

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of:

Falck-Pedersen

Application No. 08/653,114

Filed: May 24, 1996

For:

ADENOVIRUS GENE

EXPRESSION SYSTEM

Art Unit: 1632

Examiner: R. Schnizer

JAN 1 5 2002
TECH CENTER 1600/2900

PENDING CLAIMS AFTER AMENDMENTS MADE IN RESPONSE TO OFFICE ACTION DATED AUGUST 14, 2001

- 1. An adenoviral vector for expressing a heterologous gene(s) in a host cell, comprising, in an orientation opposite to the direction of adenoviral gene transcription, (a) at least one insertion site for cloning a selected heterologous gene; (b) a heterologous promoter positioned upstream from said at least one insertion site, wherein, upon cloning of the selected heterologous gene into said at least one insertion site, said gene is under the regulatory control of said heterologous promoter; (c) a eukaryotic splice acceptor and splice donor site positioned downstream of said promoter and upstream of said at least one insertion site; and (d) a polyadenylation sequence positioned downstream of said insertion site.
- 3. The adenoviral vector according to Claim 1, wherein said heterologous promoter is a mouse cytomegalovirus early promoter, or an effective expression promoting fragment thereof.
- 4. The adenoviral vector according to Claim 1, wherein said polyadenylation sequence is the mouse β -globin polyadenylation sequence.
- 9. The adenoviral vector according to Claim 1, wherein said at least one insertion site further comprises a second insertion site for insertion of a second heterologous gene.
- 17. The adenoviral vector according to Claim 1, wherein said adenoviral vector further comprises heterologous DNA inserted in said at least one insertion site.
 - 18. A host cell infected with the adenoviral vector of Claim 17.

In re Appln. of Falck-Pedersen Application No. 08/653,114

- 19. A method for producing a selected protein, comprising culturing a host cell which has been infected with the adenoviral vector of Claim 17, wherein said heterologous DNA encodes a selected protein, whereupon said selected protein is produced.
- 20. A method of delivering a heterologous gene to an animal heart *in vivo*, wherein the method comprises administering to the animal heart an adenoviral vector comprising, in an orientation opposite to the direction of adenoviral gene transcription, (a) a heterologous gene; (b) a promoter positioned upstream from the heterologous gene, the heterologous gene being under the regulatory control of the promoter; (c) a eukaryotic splice acceptor and donor site positioned downstream of the promoter and upstream of the heterologous gene; and (d) a polyadenylation sequence.